

EDWARDS'S
BOTANICAL REGISTER:

OR,

ORNAMENTAL FLOWER-GARDEN
AND SHRUBBERY:

CONSISTING OF

COLOURED FIGURES OF PLANTS AND SHRUBS,
CULTIVATED IN BRITISH GARDENS;

ACCOMPANIED BY THEIR

History, Best Method of Treatment in Cultivation, Propagation, &c.

AND

MONTHLY CHRONICLE

OF

LIBRARY
NEW YORK
BOTANICAL
GARDEN

BOTANICAL AND HORTICULTURAL NEWS.

CONTINUED

By JOHN LINDLEY, Ph. D. F.R.S. AND L.S.

PROFESSOR OF BOTANY IN UNIVERSITY COLLEGE, LONDON,

AND THE ROYAL INSTITUTION OF GREAT BRITAIN,

VICE-SECRETARY OF THE HORTICULTURAL SOCIETY,

&c. &c. &c.

Vol. 27, 1841

No! No!

1840.

OR VOL. XXVI. OF THE ENTIRE WORK.

OR VOL. XIII. OF THE NEW SERIES.

—viret semper—nec fronde caduca
Carpitur.


LONDON:

JAMES RIDGWAY, PICCADILLY.

M.DCCC.XI.

CONSERVATOIRE
BOTANIQUE

DUPLICATA DE LA BIBLIOTHEQUE
CONSERVATOIRE BOTANIQUE
VENDU



MISCELLANEOUS NOTICES.

1. PLEUROTHĀLLĪS *recurvā*; folio obovato coriaceo subobliquo petiolo pluries longiore, spicā brevi recurvā, floribus pubescentibus, sepalis lateralibus connatis, petalis membranaceis rhombeis acutis, labello membranaceo ovato utrinque sinuato obtuso, columnæ margine petaloideo incurvo.

A small creeping plant, with short recurved spikes of dull pale purple flowers. Communicated by Mr. Geo. Graves from the garden of Mrs. Cannon of Stratford.

2. PLEUROTHĀLLĪS *luteolā*; folio ovato canaliculato acuto tridentato petiolo suo breviorē, pedunculo trifloro folio adpresso eoque breviorē, floribus glabris, sepalis lateralibus connatis omnibus carinatis, petalis spathulatis emarginatis, labello lineari obtuso canaliculato glabro.

Another new plant from the same collection as the last, with small yellow flowers. Mr. Graves informs me that both exhale the smell of a sliced cucumber in the middle of the day.

3. APÖRŪM *sinuatūm*; foliis lanceolatis æquilateris approximatis acutis, floribus solitariis axillaribus, labello cuneato elongato intra apicem lineā hippocrepicā crassā sinuatā circumdato.

From Sincapore, where it was collected by Mr. Cuming. It has the habit of *A. anceps*, but its leaves are much narrower and longer, and the flowers are pale yellowish green. Communicated by Messrs. Loddiges, (132).

4. GONGÖRĀ *fulvā*; var. *vitellina*.

There is a very pretty Gongora in several collections, with bright yellow flowers less spotted than usual, and altogether of a more lively aspect. It is from Mexico, and differs from *G. fulva* in having smaller leaves, and the upper half of the labellum distinctly shorter than the lower. It however

108. BOSSLĚĀ *paucifoliā*; glabra, inermis, ramis compressis alatis, foliis paucis obovatis oblongis linearibusve, pedicellis folio brevioribus, legumine glabro latitudine pluries longiore. *Bentham mss.*

A little Swan River bush, with winged branches, a few scattered obovate leaves, and yellow and brown flowers of by no means a striking appearance. I have been favoured with specimens by R. Mangles, Esq. of Sunning Hill, and by Messrs. Lowe and Co. of Clapton, and it has also flowered in the garden of the Horticultural Society.

- ✓ 109. EPIDĚNDRŪM (Euepidendrum?) *lacertinūm*; floribus racemosis, ovarii subsecundis longissimis pendulis bracteis setaceis multoties longioribus, sepalis lanceolatis acuminatis, petalis lineari-lanceolatis, labelli adnati trilobi basi bilamellati laciniis lateralibus triangularibus intermediā lineari acuminatā elongatā, antherā immersā, ovario cuniculato.

A singular plant from Guatemala, for which I am obliged to Mr. Bateman. I have only seen the flowers; but as it is stated to have the habit of *Galeandra Baueri* it probably belongs to the section of this genus which I have called *Euepidendrum*. The flowers hang down on long stalk-like ovaries from one side of a short raceme. They are bright green with the exception of the column, which is yellow, and the lip which is a little stained with purple, and has very much the appearance of a lizard's tail, the hind legs being outside the flower, and the head and shoulders buried within the cup.

110. CYPRĪPĚDIŪM (5. *Acaulia*) *barbatūm*; acaule, foliis coriaceis acutis canaliculatis maculatis, scapo pubescente, sepalo dorsali cuspidato obtuso ciliato margine revoluto, petalis lineari-oblongis subundulatis fimbriatis margine superiore verrucoso, stamine sterili lunato pubescente.

A very fine herbaceous plant, allied to *C. venustum*, *insigne*, and *purpuratum*, gathered at Singapore, on Mount Ophir by Mr. Cuming. It has flowers richly stained with purple, and streaked with green veins upon a white ground. It has flowered with Messrs. Loddiges.

111. ORNITHŌGĀLŪM (*Chlorogalum*) *divaricatūm*; foliis glaucis longissimis canaliculatis humifusis, paniculā divaricatā, perianthii laciniis linearibus undulatis apice coherentibus mox patentibus, filamentis subulatis apice incurvis, ovarii loculis dispermis.